

CAUTION

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING



Primo[®] HG

Turf Growth Regulator

syngenta[®]

ACTIVE CONSTITUENT: 120 g/L TRINEXAPAC-ETHYL

For the regulation of leaf and stem growth of grass species and as an aid in turf management as per the Directions for Use

Syngenta Australia Pty Ltd

Level 1, 2 Lyonpark Road, Macquarie Park NSW 2113

In a transport emergency dial 000, Police or Fire Brigade

For specialist advice in an emergency only, call 1800 033 111 (24 hours)

APVMA Approval No: 94862/143884

[®]

HOW TO USE

DO NOT apply to lawns or turf recovering from wear, renovation or turf affected by disease, waterlogging or other stresses (including turf going in or out of dormancy).

DO NOT use where growth regulatory effects are likely following the application of a fungicide or any other material.

DO NOT apply if rainfall is expected within 1 hour.

DO NOT water or irrigate lawns or turf for at least 1 hour after application.

DO NOT apply to Kentucky Bluegrass as damage may occur.

PRIMO HG is a lawn growth regulator which can reduce leaf and stem elongation and speed of growth, leading to reducing mowing frequency, creating a greener lawn and developing more resilient and denser lawns. Recommended applications rates are designed to give approximately a 20 to 50% reduction in clipping yield over a 2 to 6 week period. Growth regulation begins 3 to 5 days after application. The amount of growth regulation achieved will vary between situations due to environmental conditions and lawn management practices. Therefore, application rates will need to be adjusted to match growing conditions, management practices and the amount of growth regulation required in each individual situation.

Situation	Grass type	Rate	Comments
Established lawns and turf grass	Buffalo grass, Common couch, Hybrid couch eg Greenlees Park, Santa Ann, Wintergreen	Use 10 – 40 mL per 10 L of water and spray over 100 m ²	For best results apply PRIMO HG when lawns are actively growing, well maintained and receive regular water and fertiliser applications. Use the lower rates when grass growth is moderate.
	Kikuyu	Use 20 – 80 mL per 10 L of water and spray over 100 m ²	Use higher rates when grass growth is vigorous or when longer periods of 4 to 6 weeks regulation are required.
	Fescue grass, Perennial rye grass	Use 20 – 60 mL per 10 L of water and spray over 100 m ²	Repeat applications can be made as soon as turf resumes growth or if more suppression is required.
	Carpet grass, Bent grass, Bahia grass	Use 20 – 40 mL per 10 L water and spray over 100 m ²	Apply as part of a program to maintain ongoing regulation of growth.
	Winter grass reduction (in Bent grass)	Use 4 – 10 mL per 10 L water and spray over 100 m ²	Winter Grass Management in Bent Grass. Apply at peak growth periods. Low rates of PRIMO HG can provide growth suppression of Winter Grass, but not Bent grass. This can enable a transition towards a higher percentage of Bent grass.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIODS

DO NOT GRAZE TREATED TURF/LAWN, OR FEED TURF/LAWN CLIPPINGS FROM ANY TREATED AREA TO POULTRY OR LIVESTOCK

HOW TO PREPARE

1. Prior to opening, shake the PRIMO HG bottle well.
2. Add half the required amount of water to your sprayer.
3. Add the required quantity of PRIMO HG to the half full sprayer.
4. Add the remaining water required and ensure it is closed before shaking sprayer.
5. Only mix sufficient chemical to be used on the day of mixing for the application area.

HOW TO APPLY

- For best results do not mow 6 hours before or after application.
- Measure the lawn area before application to determine the size. Make up only enough mixture to apply over the area.
- Evenly spray the mixed solution over the area to be treated.
- Thoroughly clean all equipment after application. Empty and thoroughly rinse sprayer with clean water.

CLEAN UP

Wash out spraying equipment and mixing and measuring utensils with water onto grass.

For more information regarding application visit syngentaturf.com.au

CAUTION

DO NOT allow chemical containers or spray to get into drains, sewers, streams or ponds. DO NOT allow re-entry into treated areas until the spray has dried.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, dry place out of the reach of children. DO NOT store in direct sunlight. Dispose of empty container by wrapping in paper, placing in a plastic bag and putting in garbage.

SAFETY DIRECTIONS

May irritate eyes and skin. Avoid contact with eyes and skin. Wash hands after use.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131 126.

DISCLAIMER

This product complies with the specifications in its statutory registration. Implied terms and warranties are excluded. Syngenta's liability for breach of the express or any non-excludable implied warranty is limited to product replacement or purchase price refund. The purchaser must determine suitability for intended purpose and take all proper precautions in the handling, storage and use of the product including those on the label and/or safety data sheet failing which Syngenta shall have no liability.

Product names marked ® or ™, the ALLIANCE FRAME
the SYNGENTA Logo and the PURPOSE ICON
are Trademarks of a Syngenta Group Company

